

Draft of February 2, 2008

A GENERAL FORMULA FOR THE ALGEBRAIC DEGREE IN SEMIDEFINITE PROGRAMMING

HANS-CHRISTIAN GRAF VON BOTHMER AND KRISTIAN RANESTAD

ABSTRACT. In this note, we use a natural desingularization of the conormal variety of the variety of $(n \times n)$ -symmetric matrices of rank at most r to find a general formula for the algebraic degree in semidefinite programming.

1. THE ALGEBRAIC DEGREE IN SEMIDEFINITE PROGRAMMING

Let $\mathbb{P}^m$ be a general projective space of symmetric $(n \times n)$ -matrices up to scalar multiples, and let $Y_r \subset \mathbb{P}^m$ be the subvariety of matrices of rank at most r . In this note we find a formula for the degree $\delta(m, n, r)$ of the dual variety Y_r^* whenever this is a hypersurface, i.e. $\delta(m, n, r)$ is the number of hyperplanes in $\mathbb{P}^m$ in a general pencil that are tangent to Y_r at some smooth point.

In [3] the algebraic degree of semi-definite programming is introduced and is shown to coincide with a degree of a dual variety as above [3, Theorem 5.2]. In particular, the number $\delta(m, n, r)$ is a measure of the algebraic complexity of the rank r solution for a general objective function optimized over a generic m -dimensional affine space of symmetric $(n \times n)$ -matrices.

Our general formula for $\delta(m, n, r)$ extends the results of [3]. It is expressed in terms of a function on subsequences of $\{1, \dots, n\}$. Let

$$\psi_i = 2^{i-1}, \quad \psi_{i,j} = \sum_{k=i}^{j-1} \binom{i+j-2}{k} \quad \text{when } i < j,$$

and

$$\begin{aligned} \psi_{i_1, \dots, i_r} &= \text{Pf}(\psi_{i_k, i_l})_{1 \leq k < l \leq r} & \text{if } r \text{ is even} \\ \psi_{i_1, \dots, i_r} &= \text{Pf}(\psi_{i_k, i_l})_{0 \leq k < l \leq r} & \text{if } r \text{ is odd} \end{aligned}$$

where $\psi_{i_0, i_k} = \psi_{i_k}$ and Pf denotes the Pfaffian.

Theorem 1.1. *The algebraic degree*

$$\delta(m, n, r) = \sum_I \psi_I \psi_{I^c},$$

where the sum runs over all strictly increasing subsequences $I = \{i_1, \dots, i_{n-r}\}$ of $\{1, \dots, n\}$ of length $n - r$ and sum $i_1 + \dots + i_{n-r} = m$, and I^c is the complement $\{1, \dots, n\} \setminus I$.

Notice that the formula makes sense only when $m \geq \binom{n-r+1}{2}$ and $\binom{r+1}{2} \leq \binom{n+1}{2} - m$, which is precisely the allowable range for the algebraic degree (cf. [3, Proposition 3.1]).

2. THE BIDEGREE OF A CONORMAL VARIETY

Let V and V^* be an n -dimensional vector space and its dual with fixed dual bases. Then we interpret a symmetric $(n \times n)$ -matrix as a bilinear map $\phi_S : V \rightarrow V^*$ (or $\phi_T : V^* \rightarrow V$) such that $\phi(v_1)(v_2) = \phi(v_2)(v_1)$ for any pair of vectors v_1, v_2 . The symmetric maps $V \rightarrow V^*$ form S^2V^* , while the symmetric maps $V^* \rightarrow V$ form S^2V . In this notation we let $Y_r \subset \mathbb{P}(S^2V)$ be the space of symmetric $(n \times n)$ -matrices of rank at most r (up to scalars) and $CY_r \subset \mathbb{P}(S^2V^*) \times_{\mathbb{C}} \mathbb{P}(S^2V)$ the variety of pairs of symmetric matrices (S, T) of rank at most $n - r$ and r whose matrix product $ST = 0$. The variety CY_r is the conormal variety of Y_r , i.e. the closure of the incidence set of pairs (S, T) such that $S \in \mathbb{P}(S^2V^*)$ corresponds to a hyperplane in $\mathbb{P}(S^2V)$ that is tangent to Y_r at the smooth point T .

The bidegree of $CY_r \subset \mathbb{P}(S^2V^*) \times_{\mathbb{C}} \mathbb{P}(S^2V)$ is the generating function for the algebraic degree. More precisely, denote by H_1 be the class of a hyperplane on $\mathbb{P}(S^2V^*)$ and H_2 be the class of a hyperplane on $\mathbb{P}(S^2V)$, then the class of CY_r is a homogeneous polynomial in H_1 and H_2 .

Proposition 2.1. [3, Theorem 4.2]

$$[CY_r] = \sum_m \delta(m, n, r) H_1^{\binom{n+1}{2}-m} H_2^m$$

We may therefore compute the algebraic degree on CY_r :

Corollary 2.2.

$$\delta(m, n, r) = \int_{CY_r} H_1^{m-1} H_2^{\binom{n+1}{2}-m-1} \cap CY_r.$$

3. A DESINGULARIZATION OF THE CONORMAL VARIETY

Following the argument of [3, Lemma 3.2] we define a natural desingularization of CY_r : Fix a rank r subspace K of V . The set of pairs (S, T) of symmetric bilinear maps $S : V \rightarrow V^*$ and $T : V^* \rightarrow V$ such that $K \subset \ker S$ and $K^\perp \subset \ker T \subset V^*$ form a product of projective spaces: S determines an element in $S^2(V/K)^* \subset S^2V^*$, while T determines an element in $S^2K \subset S^2V$, since $(V^*/K^\perp)$ is naturally isomorphic to K^* .

Varying K we find a desingularization of CY_r as a fiber bundle over the Grassmannian of rank r subspaces of V . More precisely, consider the Grassmannian $G = G(r, V)$ of rank r subspaces, and the universal sequence

$$0 \rightarrow U \rightarrow V_G \rightarrow Q \rightarrow 0$$

on G . Consider the symmetric squares S^2U and S^2Q^* and their projectivisations $\mathbb{P}(S^2U)$ and $\mathbb{P}(S^2Q^*)$. The inclusions $S^2U \subset S^2V$ and $S^2Q^* \subset S^2V^*$ define a natural map

$$X := \mathbb{P}(S^2Q^*) \times_G \mathbb{P}(S^2U) \rightarrow \mathbb{P}(S^2V^*) \times_{\mathbb{C}} \mathbb{P}(S^2V).$$

Proposition 3.1. *Let $G = G(r, V)$ and $X = \mathbb{P}(S^2Q^*) \times_G \mathbb{P}(S^2U)$, where Q and U are the universal quotient and subbundle on G . Then the image of the natural projection $X \rightarrow \mathbb{P}(S^2V^*) \times_{\mathbb{C}} \mathbb{P}(S^2V)$ is the conormal variety CY_r of the variety Y_r of symmetric bilinear maps $T : V^* \rightarrow V$ of rank at most r , and $X \rightarrow CY_r$ is a desingularization of CY_r .*

Proof. On the one hand, the image of the projection consists of all pairs (S, T) , such that the composition $ST = 0$ and the ranks of S and T are at most $n - r$ and r , so the image is precisely the conormal variety CY_r of Y_r . On the other hand the variety X is clearly smooth, and the map $X \rightarrow CY_r$ is birational, so it is a desingularization of CY_r . $\square$

4. THE PROOF OF THE FORMULA

We denote by H_1 and H_2 also their pullback to X . It remains by Corollary 2.2 to compute $H_1^a H_2^b$ on X , when $a + b = \binom{n+1}{2} - 2 = r(n - r) + \binom{r+1}{2} - 1 + \binom{n-r+1}{2} - 1 = \dim X$. Let $p_1 : \mathbb{P}(S^2 Q^*) \rightarrow G$ and $p_2 : \mathbb{P}(S^2 U) \rightarrow G$ be the natural projections and $p : X \rightarrow G$ be the induced projection from the fiber product. We use Segre classes to move the computation from X to G . Our projective bundles are the varieties of rank one subbundles and not rank one quotient bundles as in [1], so the Segre classes in [1, 2.4] are the Segre classes of the dual bundles in our notation. Since $S^2 Q^*$ has rank $\binom{n-r+1}{2}$, we therefore get

$$(p_1)_*(H_1^{\binom{n-r+1}{2}-1+i} \cap X) = (-1)^i s_i S^2 Q^* \cap G = s_i S^2 Q \cap G.$$

Thus

$$(p_1)_*(H_1^a \cap X) = s_{a-\binom{n-r+1}{2}+1} S^2 Q \cap G.$$

Similarly we compute H_2^b :

$$(p_2)_*(H_2^b \cap X) = s_{b-\binom{r+1}{2}+1} S^2 U^* \cap G.$$

On the fiber product we get

$$p_*(H_1^a H_2^b \cap X) = p_*(H_1^a H_2^b \cap p^* G) = s_{a-\binom{n-r+1}{2}+1} (S^2 Q) s_{b-\binom{r+1}{2}+1} (S^2 U^*) \cap G.$$

Combined with Corollary 2.2 we have therefore reduced the computation of $\delta(m, n, r)$ to a calculation on the Grassmannian variety G .

Proposition 4.1. *The algebraic degree*

$$\delta(m, n, r) = \int_G s_{m-\binom{n-r+1}{2}} (S^2 Q) s_{\binom{n+1}{2}-m-\binom{r+1}{2}} (S^2 U^*) \cap G.$$

Proof. Since X is birational to CY_r and $H_1^{m-1} H_2^{\binom{n+1}{2}-m-1} \cap CY_r$ is finite,

$$\delta(m, n, r) = \int_{CY_r} H_1^{m-1} H_2^{\binom{n+1}{2}-m-1} \cap CY_r = \int_X H_1^{m-1} H_2^{\binom{n+1}{2}-m-1} \cap X.$$

$\square$

In this formulation, computing the degree involves Schubert calculus on G . In two steps we will circumvent this, following some basic results on Schur functions on Grassmannians (cf. [2] and [1]) : First, we express the total Segre classes of the symmetric product with the Segre classes $s_i = s_i(E)$ of the original bundle:

$$s(S^2 E) = \sum_I \psi_I s_I(E)$$

where I runs over all strictly increasing sequences $\{i_1, \dots, i_e\}$ of positive integers of length equal to the rank e of E , and $s_I(E)$ is the Schur function

$$s_I(E) = \det \begin{pmatrix} s_{i_1-1} & s_{i_2-1} & \dots & s_{i_e-1} \\ s_{i_1-2} & s_{i_2-2} & \dots & s_{i_e-2} \\ \dots & \dots & \dots & \dots \\ s_{i_1-e} & s_{i_2-e} & \dots & s_{i_e-e} \end{pmatrix}.$$

In particular $s_I(E)$ has degree $i_1 + \dots + i_e - \binom{e+1}{2}$. The coefficient ψ_I is the integral valued function

$$\psi_I = \sum_J A_{I,J},$$

where J runs over all strictly increasing sequences of length equal to the length of I , and $A_{I,J}$ is the minor of following matrix with rows indexed by I and columns indexed by J

$$A = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & \dots \\ 1 & 1 & 0 & 0 & 0 & 0 & \dots \\ 1 & 2 & 1 & 0 & 0 & 0 & \dots \\ 1 & 3 & 3 & 1 & 0 & 0 & \dots \\ 1 & 4 & 6 & 4 & 1 & 0 & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \end{pmatrix}$$

We refer to [1, Proposition A.15], for the following closed formulas for the function ψ_I :

$$\psi_i = 2^{i-1}, \quad \psi_{i,j} = \sum_{k=i}^{j-1} \binom{i+j-2}{k} \quad \text{when } i < j,$$

and

$$\begin{aligned} \psi_{i_1, \dots, i_r} &= \text{Pf}(\psi_{i_k, i_l})_{1 \leq k < l \leq r} \quad \text{if } r \text{ is even} \\ \psi_{i_1, \dots, i_r} &= \text{Pf}(\psi_{i_k, i_l})_{0 \leq k < l \leq r} \quad \text{if } r \text{ is odd} \end{aligned}$$

where $\psi_{i_0, i_k} = \psi_{i_k}$ and Pf denotes the Pfaffian.

Thus,

$$s_{m - \binom{n-r+1}{2}}(S^2 Q) = \sum_I \psi_I s_I(Q)$$

where I runs over all strictly increasing sequences of positive integers $I = \{i_1, i_2, \dots, i_{n-r}\}$ with sum $i_1 + \dots + i_{n-r} = m$. (Notice that our sequences are shifted once to the right compared to those of [1], in particular they do not contain 0.)

Secondly, if I has length $n-r$ and J has length r , then

$$s_J(U) s_I(Q) = s_{JI} V_G$$

where JI is the concatenated sequence (cf. [1, 3.1]). Now, $s_{JI} = 0$ whenever the sequence JI has repeated entries. Furthermore, since V_G is a trivial bundle, $s_{JI} V_G \cap G = 0$ whenever the degree of $s_{JI} V_G$ is positive, i.e. the sum of the entries in JI exceeds $\binom{n+1}{2}$. We conclude that $s_{JI} V_G \cap G$ is nonzero only if JI is a permutation of $\{1, \dots, n\}$. In particular, $i_{n-r}, j_r \leq n$ and, by (cf. [1, 3.2.1])

$$s_I(Q) s_J(U^*) \cap G = \delta_{IJ^c}$$

where I and J are subsequences of $\{1, \dots, n\}$ of length $n - r$ and r respectively, and the sequence J^c is the complement $\{1, \dots, n\} \setminus J$ and, finally, $\delta_{AB} = 1$ if $A^c = B$ and $\delta_{AB} = 0$ otherwise.

We put the two steps together and the Theorem 1.1 immediately follows.

Example 4.2. We compute $\delta(m, n, r)$, when $n = 5$. The relevant values of ψ are:

$$\begin{array}{c} I \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \\ \psi_I \quad 1 \quad 2 \quad 4 \quad 8 \quad 16 \end{array}$$

$$\begin{array}{c} I \quad 1, 2 \quad 1, 3 \quad 1, 4 \quad 1, 5 \quad 2, 3 \quad 2, 4 \quad 2, 5 \quad 3, 4 \quad 3, 5 \quad 4, 5 \\ \psi_I \quad 1 \quad 3 \quad 7 \quad 15 \quad 3 \quad 10 \quad 25 \quad 10 \quad 35 \quad 35 \end{array}$$

$$\begin{array}{c} I \quad 1, 2, 3 \quad 1, 2, 4 \quad 1, 2, 5 \quad 1, 3, 4 \quad 1, 3, 5 \quad 1, 4, 5 \quad 2, 3, 4 \quad 2, 3, 5 \quad 2, 4, 5 \quad 3, 4, 5 \\ \psi_I \quad 1 \quad 4 \quad 11 \quad 6 \quad 23 \quad 27 \quad 4 \quad 18 \quad 30 \quad 20 \end{array}$$

$$\begin{array}{c} I \quad 1, 2, 3, 4 \quad 1, 2, 3, 5 \quad 1, 2, 4, 5 \quad 1, 3, 4, 5 \quad 2, 3, 4, 5 \\ \psi_I \quad 1 \quad 5 \quad 10 \quad 10 \quad 5 \end{array}$$

So

$$\delta(1, 5, 4) = \psi_1 \psi_{2,3,4,5} = 5 \quad \delta(2, 5, 4) = \psi_2 \psi_{1,3,4,5} = 20$$

$$\delta(3, 5, 4) = \psi_3 \psi_{1,2,4,5} = 40 \quad \delta(4, 5, 4) = \psi_4 \psi_{1,3,4,5} = 40$$

$$\delta(5, 5, 4) = \psi_5 \psi_{1,2,3,4} = 16 \quad \delta(3, 5, 3) = \psi_{1,2} \psi_{3,4,5} = 20$$

$$\delta(4, 5, 3) = \psi_{1,3} \psi_{2,4,5} = 90$$

$$\delta(5, 5, 3) = \psi_{1,4} \psi_{2,3,5} + \psi_{2,3} \psi_{1,4,5} = 7 \cdot 18 + 3 \cdot 27 = 207$$

$$\delta(6, 5, 3) = \psi_{1,5} \psi_{2,3,4} + \psi_{2,4} \psi_{1,3,5} = 15 \cdot 4 + 10 \cdot 23 = 290$$

$$\delta(7, 5, 3) = \psi_{2,5} \psi_{1,3,4} + \psi_{3,4} \psi_{1,2,5} = 25 \cdot 6 + 10 \cdot 11 = 260$$

$$\delta(8, 5, 3) = \psi_{3,5} \psi_{1,2,4} = 140 \quad \delta(9, 5, 3) = \psi_{4,5} \psi_{1,2,3} = 35$$

Example 4.3. With a computer it takes less then one minute to compute

$$\delta(105, 20, 10) = 167223927145503062075691969268936976274880$$

REFERENCES

- [1] Laksov, D., Lascoux, A., Thorup, A. *On Giambelli's theorem on complete correlations*, Acta Math. **162**, 1989, 143-199
- [2] Lascoux, A. *Fonctions de Schur et grassmanniennes*, C.R. Acad. Sci. Paris. **281** 1975, 813-815; II, 851-854
- [3] Nie, J., Ranestad, K., Sturmfels, B. *The algebraic degree of semidefinite programming*, math.CO/0611562

INSTITUT FÜR ALGEBRAISCHE GEOMETRIE, LEIBNITZ UNIVERSITÄT HANNOVER, WELFENGARTEN 1,
D-30167 HANNOVER

E-mail address: bothmer@math.uni-hannover.de

URL: <http://www.iag.uni-hannover.de/~bothmer/>

MATEMATISK INSTITUTT, UNIVERSITETET I OSLO, PO Box 1053, BLINDERN, NO-0316 OSLO, NOR-
WAY

E-mail address: ranestad@math.uio.no

URL: <http://www.math.uio.no/~ranestad>